

## RESEARCH

## Open Access



# One Diophantine inequality with unlike powers of prime variables

Wenxu Ge<sup>1\*</sup> and Weiping Li<sup>2</sup>

\*Correspondence:

gewenxu1982@163.com

<sup>1</sup>Department of Mathematics and Information Sciences, North China University of Water Resources and Electric Power, Zhengzhou, 450046, P.R. China

Full list of author information is available at the end of the article

**Abstract**

In this paper, we show that if  $\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5$  are nonzero real numbers not all of the same sign,  $\eta$  is real,  $0 < \sigma < \frac{1}{720}$ , and at least one of the ratios  $\lambda_i/\lambda_j$  ( $1 \leq i < j \leq 5$ ) is irrational, then the inequality  $|\lambda_1 p_1 + \lambda_2 p_2^2 + \lambda_3 p_3^3 + \lambda_4 p_4^4 + \lambda_5 p_5^5 + \eta| < (\max_{1 \leq j \leq 5} p_j^j)^{-\sigma}$  has infinite solutions with primes  $p_1, p_2, p_3, p_4, p_5$ .

**MSC:** 11D75; 11P55**Keywords:** Davenport-Heilbronn method; prime; Diophantine approximation

## 1 Introduction

Diophantine inequalities with integer or prime variables have been considered by many scholars. Recently, Yang and Li in [1] proved that the inequality

$$\left| \lambda_1 x_1^2 + \lambda_2 x_2^3 + \lambda_3 x_3^4 + \lambda_4 x_4^5 - p - \frac{1}{2} \right| < \frac{1}{2}$$

has infinite solutions with natural numbers  $x_1, x_2, x_3, x_4$  and prime  $p$ . Using the Davenport-Heilbronn method, we establish our result as follows.

**Theorem 1.1** *Let  $\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5$  be nonzero real numbers not all of the same sign,  $\eta$  is real,  $0 < \sigma < \frac{1}{720}$ , and at least one of the ratios  $\lambda_i/\lambda_j$  ( $1 \leq i < j \leq 5$ ) is irrational, then the inequality*

$$|\lambda_1 p_1 + \lambda_2 p_2^2 + \lambda_3 p_3^3 + \lambda_4 p_4^4 + \lambda_5 p_5^5 + \eta| < \left( \max_{1 \leq j \leq 5} p_j^j \right)^{-\sigma}$$

*has infinite solutions with primes  $p_1, p_2, p_3, p_4, p_5$ .*

## 2 Notation and outline of the proof

Throughout, we use  $p$  to denote a prime number. We denote by  $\delta$  a sufficiently small positive number and by  $\varepsilon$  an arbitrarily small positive number, not necessarily the same at different occurrences. Constants, both explicit and implicit, in Landau or Vinogradov symbols may depend on  $\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5$ , and  $\eta$ . We write  $e(x) = e^{2\pi i x}$ . We take  $X$  to be the basic parameter, a large real integer. Since at least one of the ratios  $\lambda_i/\lambda_j$  ( $1 \leq i < j \leq 5$ ) is irrational, without loss of generality we may assume that  $\lambda_1/\lambda_2$  is irrational. For the other

cases, the only difference is in the following intermediate region, and we may deal with the same method in Section 4.

Since  $\lambda_1/\lambda_2$  is irrational, there are infinitely many pairs of integers  $q, a$  with  $|\lambda_1/\lambda_2 - a/q| \leq q^{-2}$ ,  $(a, q) = 1$ ,  $q > 0$ , and  $a \neq 0$ . We choose  $q$  to be large in terms of  $\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5, \eta$  and make the following definitions:

$$N = q^2, \quad L = \log N, \quad 0 < \sigma < \frac{\theta}{32} < \frac{1}{720}, \quad v = N^{-\sigma}, \quad \tau = N^{-1+\theta}, \quad (2.1)$$

$$P = N^\theta L^{-1}, \quad Q = (|\lambda_1|^{-1} + |\lambda_2|^{-1})N^{1-\theta}, \quad T_1 = T_2^2 = T_3^3 = T_4^4 = T_5^5 = N^{\frac{1}{3}}. \quad (2.2)$$

Let  $u$  be a positive real number, we define

$$K_u(\alpha) = \left( \frac{\sin \pi u \alpha}{\pi \alpha} \right)^2 \quad (\alpha \neq 0), \quad K_u(0) = u^2, \quad (2.3)$$

$$F_k(\alpha) = \sum_{(\delta N)^{1/k} \leq p \leq N^{1/k}} e(\lambda_k p^k \alpha) \log p, \quad k = 1, 2, 3, 4, 5, \quad (2.4)$$

$$I_k(\alpha) = \int_{(\delta N)^{1/k}}^{N^{1/k}} e(\lambda_k y^k \alpha) dy, \quad k = 1, 2, 3, 4, 5, \quad (2.5)$$

$$J_k(\alpha) = \sum_{\substack{|\gamma| \leq T_k \\ \beta \geq \frac{2}{3}}} \sum_{\delta N < n \leq N} n^{-1+\rho/k} e(\lambda_k \alpha n), \quad k = 1, 2, 3, 4, 5, \quad (2.6)$$

where  $\rho = \beta + i\gamma$  ( $\beta, \gamma$  real) is a typical non-trivial zero of the Riemann Zeta function.

It follows from (2.3) that

$$K_u(\alpha) \ll \min(u^2, |\alpha|^{-2}), \quad \int_{-\infty}^{+\infty} e(\alpha y) K_u(\alpha) d\alpha = \max(0, u - |y|). \quad (2.7)$$

From (2.7) it is clear that

$$\begin{aligned} J &:= \int_{-\infty}^{+\infty} \prod_{j=1}^5 F_j(\alpha) e(\alpha \eta) K_v(\alpha) d\alpha \\ &\leq (\log N)^5 \sum_{\substack{|\lambda_1 p_1 + \lambda_2 p_2^2 + \lambda_3 p_3^3 + \lambda_4 p_4^4 + \lambda_5 p_5^5 + \eta| < v \\ (\delta N)^{1/k} \leq p_k \leq N^{1/k}, k=1,2,3,4,5}} 1 \\ &=: (\log N)^5 \mathcal{N}(N). \end{aligned}$$

Thus we have

$$\mathcal{N}(N) \geq (\log N)^{-5} J.$$

To estimate  $J$ , we split the range of infinite integration into three sections, traditional named the neighborhood of the origin  $\mathfrak{C} = \{\alpha \in \mathbb{R} : |\alpha| \leq \tau\}$ , the intermediate region  $\mathfrak{D} = \{\alpha \in \mathbb{R} : \tau \leq |\alpha| \leq P\}$ , the trivial region  $\mathfrak{c} = \{\alpha \in \mathbb{R} : |\alpha| > P\}$ .

To prove Theorem 1.1, we shall establish that

$$J(\mathfrak{C}) \gg v^2 N^{\frac{77}{60}}, \quad J(\mathfrak{D}) = o(v^2 N^{\frac{77}{60}}), \quad J(\mathfrak{c}) = o(v^2 N^{\frac{77}{60}})$$

in Sections 3, 4, and 5, respectively. Thus

$$\mathcal{N}(N) \gg v^2 (\log N)^{-5} N^{\frac{77}{60}},$$

and Theorem 1.1 can be established.

### 3 The neighborhood of the origin

We let

$$B_k(\alpha) = F_k(\alpha) - I_k(\alpha) + J_k(\alpha), \quad k = 1, 2, 3, 4, 5. \quad (3.1)$$

We use  $C$  to denote a positive absolute constant, not necessarily the same one on each occurrence.

**Lemma 3.1** *We have*

$$B_k(\alpha) \ll N^{\frac{2}{3k}} L^C (1 + |\alpha|N), \quad k = 1, 2, 3, 4, 5. \quad (3.2)$$

*This is Lemma 7 of Vaughan [2].*

**Lemma 3.2** *For  $k = 1, 2, 3, 4, 5$ , we have*

$$I_k(\alpha) \ll N^{\frac{1}{k}} \min(1, N^{-1} |\alpha|^{-1}), \quad (3.3)$$

$$\int_{-\frac{1}{2}}^{\frac{1}{2}} |J_k(\alpha)|^2 d\alpha \ll N^{\frac{2}{k}-1} \exp(-2L^{-\frac{1}{5}}), \quad (3.4)$$

$$\int_{-\frac{1}{2}}^{\frac{1}{2}} |I_k(\alpha)|^2 d\alpha \ll N^{\frac{2}{k}-1}, \quad (3.5)$$

$$\int_{-\tau}^{\tau} |B_k(\alpha)|^2 d\alpha \ll N^{\frac{2}{k}-1} \exp(-2L^{-\frac{1}{5}}), \quad (3.6)$$

$$\int_{-\tau}^{\tau} |F_k(\alpha)|^2 d\alpha \ll N^{\frac{2}{k}-1}. \quad (3.7)$$

*Proof* The inequality (3.6) follows from (2.1) and Lemma 3.1. The others are similar to Lemma 8 of Vaughan [2].  $\square$

**Lemma 3.3** *We have*

$$\int_{\mathfrak{C}} \left| \prod_{i=1}^5 F_i(\alpha) - \prod_{i=1}^5 I_i(\alpha) \right| K_v(\alpha) d\alpha \ll v^2 N^{\frac{77}{60}} \exp(-L^{-\frac{1}{5}}). \quad (3.8)$$

*Proof* Note that

$$\begin{aligned} & \prod_{i=1}^5 F_i(\alpha) - \prod_{i=1}^5 I_i(\alpha) \\ &= (F_1(\alpha) - I_1(\alpha)) \prod_{i=2}^5 F_i(\alpha) + I_1(\alpha) (F_2(\alpha) - I_2(\alpha)) \prod_{i=3}^5 F_i(\alpha) \end{aligned}$$

$$\begin{aligned}
 & + I_1(\alpha)I_2(\alpha)(F_3(\alpha) - I_3(\alpha))F_4(\alpha)F_5(\alpha) + \prod_{i=1}^3 I_i(\alpha)(F_4(\alpha) - I_4(\alpha))F_5(\alpha) \\
 & + \prod_{i=1}^4 I_i(\alpha)(F_5(\alpha) - I_5(\alpha)).
 \end{aligned}$$

Then by (2.7), (3.1), Lemma 3.2,

$$\begin{aligned}
 & \int_{\mathbb{C}} \left| (F_1(\alpha) - I_1(\alpha)) \prod_{i=2}^5 F_i(\alpha) \right| K_v(\alpha) d\alpha \\
 & \ll \nu^2 N^{\frac{47}{60}} \int_{-\tau}^{\tau} |(B_1(\alpha) - J_1(\alpha))F_2(\alpha)| d\alpha \\
 & \ll \nu^2 N^{\frac{47}{60}} \left( \int_{-\tau}^{\tau} |(B_1(\alpha) - J_1(\alpha))|^2 d\alpha \right)^{\frac{1}{2}} \left( \int_{-\tau}^{\tau} |F_2(\alpha)|^2 d\alpha \right)^{\frac{1}{2}} \\
 & \ll \nu^2 N^{\frac{47}{60}} \left( \int_{-\tau}^{\tau} (|B_1(\alpha)|^2 + |J_1(\alpha)|^2) d\alpha \right)^{\frac{1}{2}} \\
 & \ll \nu^2 N^{\frac{77}{60}} \exp(-L^{-\frac{1}{5}}).
 \end{aligned}$$

The other cases are similar, and the proof of Lemma 3.3 is completed.  $\square$

**Lemma 3.4** *We have*

$$\int_{|\alpha| > \tau} \left| \prod_{i=1}^5 I_i(\alpha) \right| K_v(\alpha) d\alpha \ll \nu^2 N^{\frac{77}{60} - 4\theta}. \quad (3.9)$$

It follows from (2.7) and (3.3).

**Lemma 3.5** *We have*

$$\int_{-\infty}^{+\infty} \prod_{j=1}^5 I_j(\alpha) e(\alpha \eta) K_v(\alpha) d\alpha \gg \nu^2 N^{\frac{77}{60}}. \quad (3.10)$$

*Proof* To prove (3.10), we write the left side as

$$\int_{\delta N}^N \int_{(\delta N)^{\frac{1}{2}}}^{N^{\frac{1}{2}}} \cdots \int_{(\delta N)^{\frac{1}{5}}}^{N^{\frac{1}{5}}} \int_{-\infty}^{+\infty} e\left(\alpha \left(\eta + \sum_{j=1}^5 \lambda_j y_j^j\right)\right) K_v(\alpha) d\alpha dy_1 dy_2 \cdots dy_5,$$

which, by (2.7), is

$$\int_{\delta N}^N \int_{(\delta N)^{\frac{1}{2}}}^{N^{\frac{1}{2}}} \cdots \int_{(\delta N)^{\frac{1}{5}}}^{N^{\frac{1}{5}}} \max\left(0, \nu - \left|\eta + \sum_{j=1}^5 \lambda_j y_j^j\right|\right) dy_1 dy_2 \cdots dy_5. \quad (3.11)$$

We let  $z_k = y_k^k$ ,  $k = 1, 2, 3, 4, 5$ , then the integral (3.11) can be written as

$$\frac{1}{120} \int_{\delta N}^N \cdots \int_{\delta N}^N z_2^{-\frac{1}{2}} z_3^{-\frac{2}{3}} z_4^{-\frac{3}{4}} z_5^{-\frac{4}{5}} \max\left(0, \nu - \left|\eta + \sum_{j=1}^5 \lambda_j z_j\right|\right) dz_1 \cdots dz_5. \quad (3.12)$$

Since  $\lambda_1, \lambda_2, \lambda_3, \lambda_4$ , and  $\lambda_5$  are not all of the same sign, we may assume without loss of generality that  $\lambda_1 < 0, \lambda_2 > 0$ . Consider the region

$$\mathcal{B} = \{(z_2, z_3, z_4, z_5) : \delta^{\frac{1}{2}}N \leq z_2 \leq 2\delta^{\frac{1}{2}}N, \delta N \leq z_j \leq 2\delta N \ (j = 3, 4, 5)\}.$$

Then, for  $\delta$  sufficiently small and large  $N$ , whenever  $(z_2, z_3, z_4, z_5) \in \mathcal{B}$  one has

$$2\delta N < -(\lambda_2 z_2 + \lambda_3 z_3 + \lambda_4 z_4 + \lambda_5 z_5)\lambda_1^{-1} < \frac{1}{2}N$$

and so every  $z_1$  with  $|\lambda_1 z_1 + \dots + \lambda_5 z_5 + \eta| \leq \frac{1}{2}\nu$  satisfies  $\delta N < z_1 < N$ . Therefore the integral (3.12) is greater than

$$\frac{1}{480}\nu^2 \int_{\mathcal{B}} z_2^{-\frac{1}{2}} z_3^{-\frac{2}{3}} z_4^{-\frac{3}{4}} z_5^{-\frac{4}{5}} dz_2 dz_3 dz_4 dz_5 \gg \nu^2 N^{\frac{77}{60}}.$$

This completes the proof of Lemma 3.5.  $\square$

Together with Lemmas 3.3, 3.4, 3.5, we have

$$J(\mathcal{C}) = \int_{\mathcal{C}} \prod_{j=1}^5 F_j(\alpha) e(\alpha\eta) K_\nu(\alpha) d\alpha \gg \nu^2 N^{\frac{77}{60}}. \quad (3.13)$$

#### 4 The intermediate region

**Lemma 4.1** *We have*

$$\int_{-\infty}^{+\infty} |F_j(\alpha)|^{2j} K_\nu(\alpha) d\alpha \ll N^{\frac{2j}{j}-1+\varepsilon}, \quad j = 2, 3, 4, 5, \quad (4.1)$$

$$\int_{-\infty}^{+\infty} |F_1(\alpha)|^2 K_\nu(\alpha) d\alpha \ll NL. \quad (4.2)$$

*Proof* By (2.7), we have

$$\begin{aligned} & \int_{-\infty}^{+\infty} |F_2(\alpha)|^4 K_\nu(\alpha) d\alpha \\ &= \sum_{(\delta N)^{\frac{1}{2}} \leq p_1, p_2, p_3, p_4 \leq N^{\frac{1}{2}}} \prod_{i=1}^4 \log p_i \max(0, \nu - |\lambda_2(p_1^2 + p_2^2 - p_3^2 - p_4^2)|) \\ &\ll L^4 \sum_{(\delta N)^{\frac{1}{2}} \leq p_1, p_2, p_3, p_4 \leq N^{\frac{1}{2}}} \max(0, \nu - |\lambda_2(p_1^2 + p_2^2 - p_3^2 - p_4^2)|). \end{aligned}$$

Since  $N$  is large,  $|\lambda_2(p_1^2 + p_2^2 - p_3^2 - p_4^2)| < \nu$  if and only if  $p_1^2 + p_2^2 = p_3^2 + p_4^2$ . Thus, by Hua's inequality,

$$\int_{-\infty}^{+\infty} |F_2(\alpha)|^4 K_\nu(\alpha) d\alpha \ll \nu N^{1+\varepsilon}.$$

The proofs of the cases  $j = 3, 4, 5$  and (4.2) are similar.  $\square$

**Lemma 4.2** *We have*

$$\int_{-\infty}^{+\infty} |F_2(\alpha)|^2 |F_4(\alpha)|^4 K_v(\alpha) d\alpha \ll \nu N^{1+\varepsilon}. \quad (4.3)$$

*Proof* By (2.7), we have

$$\begin{aligned} & \int_{-\infty}^{+\infty} |F_2(\alpha)|^2 |F_4(\alpha)|^4 K_v(\alpha) d\alpha \\ & \ll L^6 \sum_{\substack{(\delta N)^{\frac{1}{2}} \leq p_1, p_2 \leq N^{\frac{1}{2}} \\ (\delta N)^{\frac{1}{4}} \leq p_3, p_4, p_5, p_6 \leq N^{\frac{1}{4}}}} \max(0, \nu - |\lambda_2(p_1^2 - p_2^2) - \lambda_4(p_3^4 + p_4^4 - p_5^4 - p_6^4)|) \\ & \ll \nu L^6 R(N), \end{aligned}$$

where  $R(N)$  is the number of the solutions of the equation

$$\begin{aligned} \lambda_2(p_1^2 - p_2^2) &= \lambda_4(p_3^4 + p_4^4 - p_5^4 - p_6^4), \\ (\delta N)^{\frac{1}{2}} \leq p_1, p_2 \leq N^{\frac{1}{2}}, \quad (\delta N)^{\frac{1}{4}} \leq p_3, p_4, p_5, p_6 \leq N^{\frac{1}{4}}. \end{aligned}$$

Then we have

$$R(N) \ll N^{\frac{1}{2}} \sum_{\substack{(\delta N)^{\frac{1}{4}} \leq p_3, p_4, p_5, p_6 \leq N^{\frac{1}{4}} \\ p_3^4 + p_4^4 - p_5^4 - p_6^4 = 0}} 1 + \sum_{\substack{(\delta N)^{\frac{1}{4}} \leq p_3, p_4, p_5, p_6 \leq N^{\frac{1}{4}} \\ p_3^4 + p_4^4 - p_5^4 - p_6^4 \neq 0}} d(|p_3^4 + p_4^4 - p_5^4 - p_6^4|),$$

where  $d(n)$  is the divisor function. Now (4.3) follows from [3], (2.1).  $\square$

**Lemma 4.3** ([4]) *Suppose that  $(a, q) = 1$ ,  $|\alpha - a/q| \leq q^{-2}$ , then*

$$\sum_{1 \leq p \leq X} (\log p) e(p\alpha) \ll (\log X)^5 (X^{1/2} q^{1/2} + X^{4/5} + X q^{-1/2}).$$

**Lemma 4.4** ([5]) *Suppose that  $(a, q) = 1$ ,  $|\alpha - a/q| \leq q^{-2}$ ,  $\phi(x) = \alpha x^k + \alpha_1 x^{k-1} + \dots + \alpha_{k-1} x + \alpha_k$  ( $k \geq 2$ ), then*

$$\sum_{1 \leq p \leq X} (\log p) e(\phi(p)) \ll X^{1+\varepsilon} (q^{-1} + X^{-1/2} + q X^{-k})^{4^{1-k}}.$$

**Lemma 4.5** *For  $\tau < |\alpha| \leq P$ , we have*

$$V(\alpha) := \min(F_1(\alpha), F_2(\alpha)^2) \ll N^{1-\frac{\theta}{2}+\varepsilon}.$$

*Proof* Let  $\tau < |\alpha| \leq P$ , we choose  $a_j, q_j$  ( $j = 1, 2$ ) so that  $|\lambda_j \alpha - a_j/q_j| \leq Q^{-1} q_j^{-1}$  with  $(a_j, q_j) = 1$  and  $1 \leq q_j \leq Q$ . By the method of Davenport and Heilbronn (see Lemma 11 of [6]), we have  $\max(q_1, q_2) \geq P$ . Then Lemma 4.5 follows from Lemmas 4.3 and 4.4.  $\square$

**Lemma 4.6** *We have*

$$J(\mathfrak{D}) = \int_{\mathfrak{D}} \prod_{j=1}^5 F_j(\alpha) e(\alpha \eta) K_v(\alpha) \, d\alpha \ll v^2 N^{\frac{77}{60} - (\frac{\theta}{32} - \sigma) + \varepsilon}. \quad (4.4)$$

*Proof* By Lemmas 4.1, 4.2, 4.5, and Hölder's inequality, we have

$$\begin{aligned} & \int_{\mathfrak{D}} \left| \prod_{j=1}^5 F_j(\alpha) e(\alpha \eta) K_v(\alpha) \right| \, d\alpha \\ & \ll V(\alpha)^{\frac{1}{16}} \int_{-\infty}^{+\infty} \left| (F_1(\alpha)^{\frac{15}{16}} F_2(\alpha) + F_1(\alpha) F_2(\alpha)^{\frac{7}{8}}) \prod_{j=3}^5 F_j(\alpha) \right| K_v(\alpha) \, d\alpha \\ & \ll V(\alpha)^{\frac{1}{16}} \left( \int_{-\infty}^{+\infty} |F_1(\alpha)|^2 K_v(\alpha) \, d\alpha \right)^{\frac{15}{32}} \left( \int_{-\infty}^{+\infty} |F_2(\alpha)|^4 K_v(\alpha) \, d\alpha \right)^{\frac{1}{8}} \\ & \quad \times \left( \int_{-\infty}^{+\infty} |F_2(\alpha)|^2 F_4(\alpha)^4 |K_v(\alpha)| \, d\alpha \right)^{\frac{1}{4}} \left( \int_{-\infty}^{+\infty} |F_3(\alpha)|^8 K_v(\alpha) \, d\alpha \right)^{\frac{1}{8}} \\ & \quad \times \left( \int_{-\infty}^{+\infty} |F_5(\alpha)|^{32} K_v(\alpha) \, d\alpha \right)^{\frac{1}{32}} + V(\alpha)^{\frac{1}{16}} \left( \int_{-\infty}^{+\infty} |F_1(\alpha)|^2 K_v(\alpha) \, d\alpha \right)^{\frac{1}{2}} \\ & \quad \times \left( \int_{-\infty}^{+\infty} |F_2(\alpha)|^4 K_v(\alpha) \, d\alpha \right)^{\frac{3}{32}} \left( \int_{-\infty}^{+\infty} |F_2(\alpha)|^2 F_4(\alpha)^4 |K_v(\alpha)| \, d\alpha \right)^{\frac{1}{4}} \\ & \quad \times \left( \int_{-\infty}^{+\infty} |F_3(\alpha)|^8 K_v(\alpha) \, d\alpha \right)^{\frac{1}{8}} \left( \int_{-\infty}^{+\infty} |F_5(\alpha)|^{32} K_v(\alpha) \, d\alpha \right)^{\frac{1}{32}} \\ & \ll v N^{\frac{77}{60} - \frac{\theta}{32} + \varepsilon} \ll v^2 N^{\frac{77}{60} - (\frac{\theta}{32} - \sigma) + \varepsilon}. \quad \square \end{aligned}$$

## 5 The trivial region

**Lemma 5.1** *Let  $G(\alpha) = \sum e(\alpha f(x_1, \dots, x_m))$ , where  $f$  is any real function and the summation is over any finite set of values of  $x_1, \dots, x_m$ . Then, for any  $A > 4$ , we have*

$$\int_{|\alpha| > A} |G(\alpha)|^2 K_v(\alpha) \, d\alpha \leq \frac{16}{A} \int_{-\infty}^{+\infty} |G(\alpha)|^2 K_v(\alpha) \, d\alpha.$$

*This is Lemma 2 of [7].*

**Lemma 5.2** *We have*

$$J(\mathfrak{c}) = \int_{\mathfrak{c}} \prod_{j=1}^5 F_j(\alpha) e(\alpha \eta) K_v(\alpha) \, d\alpha \ll v^2 N^{\frac{77}{60} - (\theta - \sigma) + \varepsilon}.$$

*Proof* By Lemmas 4.1, 4.2, 5.1, and Hölder's inequality, we have

$$\begin{aligned} & \int_{\mathfrak{c}} \left| \prod_{j=1}^5 F_j(\alpha) e(\alpha \eta) K_v(\alpha) \right| \, d\alpha \\ & \ll \frac{1}{P} \int_{-\infty}^{+\infty} \prod_{j=1}^5 |F_j(\alpha)| K_v(\alpha) \, d\alpha \end{aligned}$$

$$\begin{aligned}
&\ll \frac{1}{P} \max(|F_5(\alpha)|) \left( \int_{-\infty}^{+\infty} |F_1(\alpha)|^2 K_v(\alpha) d\alpha \right)^{\frac{1}{2}} \left( \int_{-\infty}^{+\infty} |F_2(\alpha)|^4 K_v(\alpha) d\alpha \right)^{\frac{1}{8}} \\
&\quad \times \left( \int_{-\infty}^{+\infty} |F_2(\alpha)|^2 |F_4(\alpha)|^4 K_v(\alpha) d\alpha \right)^{\frac{1}{4}} \left( \int_{-\infty}^{+\infty} |F_3(\alpha)|^8 K_v(\alpha) d\alpha \right)^{\frac{1}{8}} \\
&\ll v N^{\frac{77}{60}-\theta+\varepsilon} \ll v^2 N^{\frac{77}{60}-(\theta-\sigma)+\varepsilon}.
\end{aligned}$$

□

**Competing interests**

The authors declare that they have no competing interests.

**Authors' contributions**

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

**Author details**

<sup>1</sup>Department of Mathematics and Information Sciences, North China University of Water Resources and Electric Power, Zhengzhou, 450046, P.R. China. <sup>2</sup>Department of Mathematics and Information Science, Henan University of Economics and Law, Zhengzhou, 450046, P.R. China.

**Acknowledgements**

This work is supported by the National Natural Science Foundation of China (Grant Nos. 11371122, 11471112), the Innovation Scientists and Technicians Troop Construction Projects of Henan Province (China), the Research Foundation of North China University of Water Conservancy and Electric Power (No. 201084).

Received: 18 November 2015 Accepted: 19 January 2016 Published online: 28 January 2016

**References**

1. Yang, TQ, Li, WP: One Diophantine inequality with integer and prime variables. *J. Inequal. Appl.* **2015**, 293 (2015)
2. Vaughan, RC: Diophantine approximation by prime numbers, II. *Proc. Lond. Math. Soc.* **28**, 385-401 (1974)
3. Brüdern, J, Kawada, K: Ternary problems in additive prime number theory. *Dev. Math.* **6**, 39-91 (2002)
4. Vaughan, RC: *The Hardy-Littlewood Method*. Cambridge University Press, Cambridge (1997)
5. Harman, G: Trigonometric sums over primes I. *Mathematika* **28**, 249-254 (1981)
6. Vaughan, RC: Diophantine approximation by prime numbers, I. *Proc. Lond. Math. Soc.* **28**, 373-384 (1974)
7. Davenport, H, Roth, KF: The solubility of certain Diophantine inequalities. *Mathematika* **2**, 81-96 (1955)

**Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:**

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)